



Year: 2019

A case report: a 22-year-old septic patient with central venous pO₂ of 198 mmHg

Ehlers, Ulrike Elisabeth ; Mutter, Michael ; Fahner, Peter Jurriaan ; Pfammatter, Thomas

Abstract: Background Central venous saturation and central venous pressure can be determined with central venous catheters. Therefore, the tip of the catheter should be located in the superior vena cava. The location can be monitored by electrocardiography or X-ray. The central venous pressure curve is displayed on the monitor. The reference value of central venous saturation is >70%. Venous pO₂ is normally 35-45 mmHg and central venous pressure 1-9 mmHg. Case summary We treated a 22-year-old patient with septic shock. Central venous saturation was 100% with a pO₂ of 198 mmHg. The arterial blood gas analysis was comparatively low with saturation of 98% and pO₂ of 111 mmHg. On chest X-ray, the central venous catheter tip appeared on the left side of the heart. On echocardiography, aortic positioning was not evident. On the monitor, a 'venous pressure-like' curve was seen, that did not stand in exact correlation to the electrocardiogram curve. The computed tomography (CT) image showed placement of the catheter in the upper left pulmonary vein. The patient had a partial anomalous pulmonary venous return. Discussion The C-wave of the central venous pressure curve normally occurs after the R-wave of the electrocardiogram. If C-waves appeared before R-waves, the central venous catheter placement is not central venous and must be checked. In our case, the apparent 'venous' pO₂ in blood gas examination was higher than arterial pO₂. The catheter position had to be in an oxygenated vessel proximal to the left ventricle. A vascular anomaly was a possible diagnosis and was confirmed on CT imaging.

DOI: <https://doi.org/10.1093/ehjcr/ytz211>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-181883>

Journal Article

Published Version



The following work is licensed under a Creative Commons: Attribution-NonCommercial 4.0 International (CC BY-NC 4.0) License.

Originally published at:

Ehlers, Ulrike Elisabeth; Mutter, Michael; Fahner, Peter Jurriaan; Pfammatter, Thomas (2019). A case report: a 22-year-old septic patient with central venous pO₂ of 198 mmHg. *European Heart Journal. Case Reports*, 3(4):1-5.

DOI: <https://doi.org/10.1093/ehjcr/ytz211>

A case report: a 22-year-old septic patient with central venous pO₂ of 198 mmHg

Ulrike Elisabeth Ehlers¹*, Michael Mutter², Peter Jurriaan Fahner³, and Thomas Pfammatter⁴

¹Intensive Care Unit, Cantonal Hospital Glarus, Burgstrasse 99, 8750 Glarus, Switzerland; ²Institute of Cardiology, Cantonal Hospital Glarus, Burgstrasse 99, 8750 Glarus, Switzerland; ³Institute of Radiology, Cantonal Hospital Glarus, Burgstrasse 99, 8750 Glarus, Switzerland; and ⁴Institute of Radiology, University Hospital Zurich, Rämistrasse 100, 8091 Zurich, Switzerland

Received 14 July 2019; first decision 6 August 2019; accepted 3 November 2019; online publish-ahead-of-print 27 November 2019

Background

Central venous saturation and central venous pressure can be determined with central venous catheters. Therefore, the tip of the catheter should be located in the superior vena cava. The location can be monitored by electrocardiography or X-ray. The central venous pressure curve is displayed on the monitor. The reference value of central venous saturation is >70%. Venous pO₂ is normally 35–45 mmHg and central venous pressure 1–9 mmHg.

Case summary

We treated a 22-year-old patient with septic shock. Central venous saturation was 100% with a pO₂ of 198 mmHg. The arterial blood gas analysis was comparatively low with saturation of 98% and pO₂ of 111 mmHg. On chest X-ray, the central venous catheter tip appeared on the left side of the heart. On echocardiography, aortic positioning was not evident. On the monitor, a 'venous pressure-like' curve was seen, that did not stand in exact correlation to the electrocardiogram curve. The computed tomography (CT) image showed placement of the catheter in the upper left pulmonary vein. The patient had a partial anomalous pulmonary venous return.

Discussion

The C-wave of the central venous pressure curve normally occurs after the R-wave of the electrocardiogram. If C-waves appeared before R-waves, the central venous catheter placement is not central venous and must be checked. In our case, the apparent 'venous' pO₂ in blood gas examination was higher than arterial pO₂. The catheter position had to be in an oxygenated vessel proximal to the left ventricle. A vascular anomaly was a possible diagnosis and was confirmed on CT imaging.

Keywords

Case report • Partial anomalous pulmonary venous return (PAPVR) • Central venous saturation • Venous pressure curve • Cardiac curve correlation

Learning points

- The C-wave of the central venous pressure curve should occur after the R-wave on the electrocardiogram (ECG).
- If the C-wave of a venous curve appears before the R-wave of the ECG curve, the catheter tip is not localized in the superior vena cava.
- If the apparent 'venous' pO₂ in blood gas examination is higher than the arterial pO₂, there is a suspicion that the catheter is located in an oxygenated blood vessel, such as a pulmonary vein.

* Corresponding author. Tel: +41 55 646 39 98, Email: ulrike.ehlers@ksgl.ch

Handling Editor: Richard Alexander Brown

Peer-reviewers: Cemil Izgi and Laszlo Göbölös

Compliance Editor: Rahul Mukherjee

Supplementary Material Editor: Peregrine Green

© The Author(s) 2019. Published by Oxford University Press on behalf of the European Society of Cardiology.

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is properly cited. For commercial re-use, please contact journals.permissions@oup.com

Introduction

Managing patients on intensive care units requires a fast access to the circulatory system and the ability of continuous physiological monitoring. For such purposes, central venal catheters are often inserted, for example through the jugular vein. The catheter tip should be located in the superior vena cava at the level of the right atrium. At this position, the central venous pressure and oxygenation are measured. The reference value of oxygenation is 70–80%. A value above 70% is associated with a reduction in mortality.^{1–4}

Timeline

Day 1	23:00	Emergency room	Emergency room: admission of the somnolent patient of unclear aetiology. Intubation and insertion of an arterial catheter and a central venous line
Day 2	04:00	Intensive care unit	Entry intensive care unit
	13:00	Sepsis	State deterioration to a septic shock
	14:15	VenBG	Measurement of SvO ₂ from blood of the central venous line (SvO ₂ 100%; pO ₂ 198 mmHg)
	14:30	Control	Control: confirmation of the result from 14:15 (SvO ₂ 100%; pO ₂ 200 mmHg)
	14:35	ArtBG	Repeated measurements of arterial blood gas values (SaO ₂ 98%; pO ₂ 90–111 mmHg)
	14:50	X-ray	Chest X-ray: central venous catheter tip appeared on the left side of the heart (Figure 1)
	15:15	Echo	Echocardiography: aortic positioning of the central venous line was not evident
	16:00	Electrocardiogram (ECG)	On the monitor, a ‘venous pressure-like’ curve was seen, that did not stand in exact correlation to the ECG curve
	16:30	Computed tomography	Computed tomography image showed placement of the catheter in the upper left pulmonary vein

Case presentation

We describe a 22-year-old patient with no significant past medical history, who was hospitalized with septic shock from aspiration pneumonia. A central venous line was inserted into the left jugular vein and the blood saturation was determined. The saturation was 100%,




Figure 1 Anterior-posterior chest radiograph showing central venous catheter on the left side of the heart.

pO₂ 198 mmHg (normal 30–40 mmHg) and pCO₂ 46 mmHg (normal 40–50 mmHg), respectively. The arterial blood gas analysis showed a saturation of 98%, a pO₂ of 111 mmHg (normal 75–100 mmHg) and a pCO₂ of 48 mmHg (normal 35–45 mmHg).

A chest X-ray revealed the central venous catheter tip to be on the left side of the heart, next to the aortic arch (Figure 1).

The cervical ultrasound showed that the catheter was inserted correctly in the left jugular vein. In echocardiography, the catheter tip did not appear to be positioned in the aorta. The vena cava superior could not be seen. Monitoring demonstrated a venous-like pattern with a mean pressure of 16 mmHg (Figure 2). However, the shape of the curve seemed to be ‘atypical’ for a central venous pressure curve, as the C-wave of the central venous pressure curve represents tricuspid closure. Normally C-waves occur after the R-wave of the electrocardiogram (Figure 3). In our patient, the C-waves appeared before R-waves (arrow, Figure 2). Thus, it became clear that the catheter was placed neither in a central vein nor in the left outflow tract. The catheter had to be in an oxygenated vessel proximal to the left ventricle. Computed tomography (CT) scanning demonstrated that the central venous catheter was inserted from the left jugular vein into the upper left pulmonary vein. The diagnosis of a partial anomalous pulmonary venous return (PAPVR) was made (Figures 4 and 5). The patient was able to leave the hospital after a few weeks. He was examined in detail cardiologically. The treatment was conservative. He will have a 1-year follow-up.

Discussion

Anomalous pulmonary venous return is a rare congenital malformation in which pulmonary veins fail to join the left atrium. Instead, they are connected to a systemic vein or to the right atrium directly.^{5–7} This condition may be a total anomalous pulmonary venous return concerning both lungs or a partial malformation (PAPVR), concerning




Figure 2 Electrocardiogram and venous pressure curve with partial anomalous pulmonary venous return [C-wave appears before R-wave (arrow)].




Figure 3 Normal electrocardiogram and venous pressure curve [C-wave appears after R-wave (arrow)].

only one lung or parts of one lung.^{5–8} Partial anomalous pulmonary venous return is a condition that is rarely seen in patients. It has a prevalence of 0.4–0.7%⁹ and is more common on the right side with



Figure 4 Computed tomography image partial anomalous pulmonary venous return with central venous catheter in upper left pulmonary vein.

10%.¹⁰ On the left side, PAPVR has a prevalence of 0.05%.¹¹ Total anomalous pulmonary venous return is found in approximately 0.08/1000 live births,¹² which corresponds to a relative proportion of 0.4–0.9% of all congenital heart defects.⁵

Partial anomalous pulmonary venous return is often associated with other cardiac anomalies, like persistent left superior vena cava, the most common congenital malformation of thoracic return and present in 0.3–0.5% of individuals in the general population.¹³ Partial anomalous pulmonary venous return can also come along with a persistent foramen ovale or an atrial septal defect^{5,10} or can be an isolated finding, like in our case.

Most patients with PAPVR are asymptomatic and tend to go unnoticed until adulthood.¹⁴ If identified and asymptomatic, isolated PAPVR are followed-up conservatively as there is substantial risk of stenosis after rerouting an anomalous pulmonary vein. If PAPVR shows symptoms rerouting must be considered. The morbidity is relatively low. Rerouting can be already done in childhood.⁸

Mostly, the natural history dictates that, if significant left to right shunt exists, patients may develop irreversible pulmonary hypertension, pulmonary vascular obstructive disease, or right heart failure.¹⁰

Our case is interesting, because the diagnosis PAPVR was made by central venous catheterization data and not primarily from imaging. After conspicuous blood gas analysis was performed, further examination revealed that the venous curve was not in usual correlation to the Electrocardiogram (ECG) curve (Figure 2). The diagnosis was made through CT scanning (Figures 4 and 5).





Figure 5 Three-dimensional computed tomography image partial anomalous pulmonary venous return of left upper pulmonary vein.

Take home message: If venous blood gas analysis from a central venous catheter indicates abnormal high oxygen levels, but the shape of the venous pressure curve is similar to a venous curve, a comparison with an ECG curve should be made.

Lead author biography



KD Dr Ulrike Elisabeth Ehlers, eMBA, performed her undergraduate studies at the Georg-August-University Göttingen, Germany with internships in St. Lucia (West Indies), Hobart (Australia), and Davos (Switzerland). Doctor thesis and Clinical lecturer at the University of Zurich. Specialist in Internal Medicine FMH and Intensive Care Medicine FMH, European Diploma of Intensive

Care Medicine, 'Emergency doctor SGNOR', Ultrasound specialist 'POCUS SGUM'. Executive Master in Business Administration with focus on Medical Management. Education in 'Advanced Studies in Applied Ethics' at the University of Zurich.

Supplementary material

[Supplementary material](#) is available at *European Heart Journal - Case Reports* online.

Acknowledgements

We are indebted to the employees of the Intensive Care Unit Glarus for their active contribution to the management of the patient. We thank Dr Moritz Schürch for his professional support and Dr Bert Rost for assistance in completing the 3D reconstruction.

Slide sets: A fully edited slide set detailing this case and suitable for local presentation is available online as [Supplementary data](#).

Consent: The author/s confirm that written consent for submission and publication of this case report including image(s) and associated text has been obtained from the patient in line with COPE guidance.

Conflict of interest: none declared.

References

1. Bloss F, Reinhart K. The value of central venous O₂ saturation for assessment of tissue oxygenation. *Dtsch Med Wochenschr* 2004;**129**:2601–2604.
2. Teboul JL, Hamzaoui O, Monnet X. SvO₂ to monitor resuscitation of septic patients: let's just understand the basic physiology. *Crit Care* 2011;**15**:1005.
3. Squara P. Central venous oxygenation: when physiology explains apparent discrepancies. *Crit Care* 2014;**18**:579.
4. Fresenius M, Heck M. *Repetitorium Intensivmedizin*, Vol. 4. Springer-Verlag Berlin Heidelberg; 2011. p.24.
5. Bertram H, Dewald O, Lindinger A. *Partielle und totale Lungenvenenfehlbildung*. Deutsche Gesellschaft für pädiatrische Kardiologie. Leitlinien. 2014; <http://www.kinderkardiologie.org/leitlinien> (15 November 2019).
6. Ellis FH, Kirklin JW. Anomalous Pulmonary Venous Connections. *Surgical Clinics of North America* 1955; **35**:997.

7. Senocak F, Özme S, Bilciç A, Özkutlu S, Özer S, Saraçlar M. Partial anomalous pulmonary venous return. Evaluation of 51 cases. *Jpn Heart J* 1994;**35**:43–50.
8. ElBardissi AWW, Dearani JA, Suri RM, Danielson GK. Left-sided partial anomalous pulmonary venous connections. *Ann Thorac Surg* 2008;**85**:1007–1014.
9. Wang TL, Hung HF, Lin CC, Hsiung MC, Wei J. Multiple anomalous left pulmonary venous connections detected with transthoracic echocardiography. *J Cardiothorac Surg* 2013;**8**:130.
10. Javangula K, Cole J, Cross M, Kay PH. An unusual manifestation of left partial anomalous pulmonary venous connection. *Interact CardioVasc Thorac Surg* 2010;**11**:846–848.
11. Van Meter C, LeBlanc JG, Culpepper WS, Ochsner JL. Partial anomalous pulmonary venous return. *Circulation* 1990;**82**:IV195–IV198.
12. Böhne M, Jack T, Köditz H, Beerbaum P. Kritische Herzfehler des Neugeborenen-Teil 2: Ausgewählte kritische Herzfehler. *Neonatal Scan* 2015;**4**: 61–88.
13. Tyrak KW, Holda J, Holda MK, Koziej M, Piątek K, Klimek- Piotrowska W. Persistent left superior vena cava. *Cardiovasc J Afr* 2017;**28**:e1–e4.
14. Edwin F. eComment: Left-sided partial anomalous pulmonary venous connection—should diagnosis lead to surgery? *Interact Cardiovasc Thorac Surg* 2010;**11**:847–848.